Author Index

Adler, J.E., see Lu, B., 359

Akitaya, T., see Mitsuhashi, M., 177

Angulo, J.A., Coirini, H., Ledoux, M. and Schumacher, M., Regulation by dopaminergic neurotransmission of dopamine D₂ mRNA and receptor levels in the striatum and nucleus accumbens of the rat, 1

Angulo, J.A., Printz, D., Ledoux, M. and McEwen, B.S., Isolation stress increases tyrosine hydroxylase mRNA in the locus coeruleus and midbrain and decreases proenkephalin mRNA in the striatum and nucleus accumbens, 301

Araki, T., see Zhang, J.-H., 239 Atkinson, B.G., see Teskey, G.C., 1

Badoyannis, H.C., Sharma, S.C. and Sabban, E.L., The differential effects of cell density and NGF on the expression of tyrosine hydroxylase and dopamine β -hydroxylase in PC12 cells, 79

Baker, H.J., see Claro, E., 265

Batter, D.K. and Kessler, J.A., Regionspecific regulation of preproenkephalin mRNA in cultured astrocytes, 65

Becker, C.-M., Seitanidou, T. and Triller, A., Conservation of antigenic epitopes of the inhibitory glycine receptor in rodent and goldfish CNS, 327

Berens, M.E., see Deschepper, C.F.,355 Berg, L., see Burke, W.J., 37

Bierer, L., see Perez, N., 249

Bigornia, V., see Deschepper, C.F., 355

Bing, G., Filer, D., Miller, J.C. and Stone, E.A., Noradrenergic activation of immediate early genes in rat cerebral cortex, 43

Black, I.B., see Lu, B., 359

Blaugrund, E., see Cohen, I., 181

Bloem, L.J., see Yu, L., 143

Bogerd, J., Geraerts, W.P.M., Van Heerikhuizen, H., Kerkhoven, R.M. and Joosse, J., Characterization and evolutionary aspects of a transcript encoding a neuropeptide precursor of *Lymnaea* neurons, VD1 and RPD2, 47

Brown, R.M., see Streamson Jr., C.C., 291 Buchwald, N.A., see Cohen, R.W., 27

Burke, W.J., O'Malley, K.L., Chung, H.D., Harmon, S.K., Miller, J.P. and Berg, L., Effect of pre- and postmortem variables on specific mRNA levels in human brain, 37

Buzzi, M.G., see Uhl, G.R., 133

Cain, C.J., see Clark, M., 55
Cain, D.P., see Teskey, G.C., 1
Campagnoni, A.T., see Cohen, R.W., 27
Carroll, P.D., see Howland, D.S., 345
Chao, M.V., see Gibbs, R.B., 207
Charya, S., see Perez, N., 249
Chung, H.D., see Burke, W.J., 37

Clark, M., Post, R.M., Weiss, S.R.B., Cain, C.J. and Nakajima, T., Regional expression of c-fos mRNA in rat brain during the evolution of amygdala kindled seizures, 55

Claro, E., Wallace, M.A., Fain, J.N., Nair, B.G., Patel, T.B., Shanker, G. and Baker, H.J., Altered phosphoinositide-specific phospholipase C and adenylyl cyclase in brain cortical membranes of cats with GM₁ and GM₂ gangliosidosis, 265

Cohen, I., Shani, Y., Blaugrund, E. and Schwartz, M., Isolation and sequence analysis of two intermediate filament cDNA clones from fish optic nerve, 181

Cohen, R.W., Fisher, R.S., Duong, T., Handley, V.W., Campagnoni, A.T., Hull, C.D., Buchwald, N.A. and Levine, M.S., Altered excitatory amino acid function and morphology of the cerebellum of the spastic Han-Wistar rat, 27

Coirini, H., see Angulo, J.A., 1 Coombs, D.W., see DeLeo, J.A., 167 Covault, J., Liu, Q.-y. and El-Deeb, S.,

Covault, J., Liu, Q.-y. and El-Deeb, S., Calcium-activated proteolysis of intracellular domains in the cell adhesion molecules NCAM and N-cadherin, 11

Davidson, N., see Yu, L., 143

DeGennaro, L.J., see Howland, D.S., 345 DeLeo, J.A., Coombs, D.W. and McCarthy, L.E., Differential c-fos-like protein expression in mechanically versus chemically induced visceral nociception, 167

Deschepper, C.F., Bigornia, V., Berens, M.E. and Lapointe, M.C., Production of thrombin and antithrombin III by brain and astroglial cell cultures, 355

Dingledine, R., see Kleckner, N.W., 1 Dionne, C.A., see Powell, P.P., 71

Donnelly-Roberts, D.L. and Lentz, T.L., Structural and conformational similarity between synthetic peptides of curaremimetic neurotoxins and rabies virus glycoprotein, 107

Dreyfus, C.F., see Lu, B., 359

Duong, T., see Cohen, R.W., 27
D'Mello, S.R. and Heinrich, G., Structural
and functional identification of regulatory regions and cis elements surrounding the nerve growth factor gene pro-

moter, 255

Eastham, H., see Whiting, P., 1 Eckermann, C.E., see Toews, A.D., 321 El-Deeb, S., see Covault, J., 11 Elferink, L.A., see Volknandt, W., 283 Elliott, R., see Lu, B., 359 Estes, P.S., see Howland, D.S., 345

Fain, J.N., see Claro, E., 265 Federoff, H.J., see Lustig, R.H., 125 Filer, D., see Bing, G., 43 Finch, C.E., see Poirier, J., 97 Finklestein, S.P., see Powell, P.P., 71 Fisher, R.S., see Cohen, R.W., 27

Gehle, V.M. and Sumikawa, K., Sitedirected mutagenesis of the conserved N-glycosylation site on the nicotinic acetylcholine receptor subunits, 17

Geraerts, W.P.M., see Bogerd, J., 47 Gibbs, R.B., Chao, M.V. and Pfaff, D.W., Effects of fimbria-fornix and angular bundle transection on expression of the p75^{NGFR} mRNA by cells in the medial septum and diagonal band of Broca: correlations with cell survival, synaptic reorganization and sprouting, 207

Handley, V.W., see Cohen, R.W., 27
Harmon, S.K., see Burke, W.J., 37
Haroutunian, V., see Perez, N., 249
Heavens, R.P., see Matthews, S.G., 171
Hein, S., see Nualart, F., 227
Heinrich, G., see D'Mello, S.R., 255
Hemmendinger, L.M., see Howland, D.S., 345

Gong, Q., see Huang, Z., 309

Hennessey, K.L., see Streamson Jr., C.C., 291

Hess, M., see Poirier, J., 97 Hirsch, J., See Streamson Jr., C.C.,291 Hoffman, B.J., see Yu, L., 143

Hosford, D.A., see Simonato, M., 115
Howland, D.S., Hemmendinger, L.M.,
Carroll, P.D., Estes, P.S., Melloni Jr.,
R.H. and DeGennaro, L.J., Positiveand negative-acting promoter sequences
regulate cell type-specific expression of
the rat synapsin I gene, 345

Huang, Z., Thewke, D., Gong, Q., Schlichter, D. and Wicks, W.D., Functional recognition of the neuronal tyrosine hydroxylase cAMP regulatory element in different cell types, 309

Hull, C.D., see Cohen, R.W., 27

Jaye, M., see Powell, P.P., 71 Johnson, G., see Perez, N., 249 Joosse, J., see Bogerd, J., 47

Kerkhoven, R.M., see Bogerd, J., 47
Kessler, J.A., see Batter, D.K., 65
Kim, J., see ?Streamson Jr., C.C.,291
Klagsbrun, M., see Powell, P.P., 71
Kleckner, N.W. and Dingledine, R., Regulation of hippocampal NMDA receptors by magnesium and glycine during development, 1.

Kozak, C.A., see Yu, L., 143 Kubo, T., see Saika, T., 187

Labiner, D.M., see Simonato, M., 115 Lapointe, M.C., see Deschepper, C.F., 355 Le, H., see Yu, L., 143 Ledoux, M., see Angulo, J.A., 1

Ledoux, M., see Angulo, J.A., 301 Lee, S.Y., see Toews, A.D., 321 Lee, J.-M., see Lu, B., 359 Leibel, R.L., see Streamson Jr., C.C.,291 Lentz, T.L., see Donnelly-Roberts, D.L., 107 Lester, H.A., see Yu, L., 143 Levine, M.S., see Cohen, R.W., 27 Liu, Q.-y., see Covault, J., 11 Loh, Y.P., see Wong, M., 197 Lu, B., Lee, J.-M., Elliott, R., Dreyfus, C.F., Adler, J.E. and Black, I.B., Regulation of NGF gene expression in CNS glia by cell-cell contact, 359 Lübbert, H., see Yu, L., 143

Lustig, R.H., Sudol, M., Pfaff, D.W. and Federoff, H.J., Estrogenic regulation and sex dimorphism of growthassociated protein 43 kDa (GAP-43) messenger RNA in the rat, 125

Mansbach, H.M., see Simoinato, M.,115 Maren, S., see Schreiber, S.S., 89 Maroteaux, L. and Scheller, R.H., The rat brain synucleins; family of proteins transiently associated with neuronal membrane, 335

Matsunaga, T., see Saika, T., 187 Matthews, S.G., Heavens, R.P. and Sirinathsinghji, D.J.S., Cellular localization of corticotropin releasing factor mRNA in the ovine brain, 171

May, P.C., see Poirier, J., 97 McCarthy, L.E., see DeLeo, J.A., 167 McEwen, B.S., see Angulo, J.A., 301 McNamara, J.O., see Simoinato, M.,115 Melloni Jr., R.H., see Howland, D.S., 345 Merril, C., see Perez, N., 249 Miller, J.C., see Bing, G., 43 Miller, J.P., see Burke, W.J., 37 Mitsuhashi, M., Akitaya, T., Turk, C.W. and Payan, D.G., Amyloid β protein substituent peptides do not interact with the substance P receptor expressed in cultured cells, 177

Morell, P., see Toews, A.D., 321 Moskowitz, M.A., see Uhl, G.R., 133 Mozley, L., see Whiting, P., 1

Nadi, N.S., see Shinoda, H., 221 Nair, B.G., see Claro, E., 265 Nakajima, T., see Clark, M., 55 Nava, S., see Whiting, P., 1 Nguyen, H., see Yu, L., 143 Nishimori, T., see Uhl, G.R., 133 Noguchi, K., see Saika, T., 187 Nualart, F., Hein, S., Rodríguez, E.M. and Oksche, A., Identification and partial characterization of the secretory glycoproteins of the bovine subcommissural organ-Reissner's fiber complex. Evidence for the existence of two precursor forms, 227

Oksche, A., see Nualart, F., 227 O'Malley, K.L., see Burke, W.J., 37

Patel, T.B., see Claro, E., 265 Payan, D.G., see Mitsuhashi, M., 177 Perez, N., Sugar, J., Charya, S., Johnson, G., Merril, C., Bierer, L., Perl, D., Haroutunian, V.and Wallace, W., Increased synthesis and accumulation of heat shock 70 proteins in Alzheimer's disease, 249

Perl, D., see Perez, N., 249 Pevsner, J., see Volknandt, W., 283 Pfaff, D.W., see Gibbs, R.B., 207 Pfaff, D.W., see Lustig, R.H., 125 Poat, J., see Whiting, P., 1 Poirier, J., Hess, M., May, P.C. and Finch,

C.E., Astrocytic apolipoprotein E mRNA and GFAP mRNA in hippocampus after entorhinal cortex lesioning, 97 Post, R.M., see Clark, M., 55

Powell, P.P., Finklestein, S.P., Dionne, C.A., Jaye, M. and Klagsbrun, M., Temporal, differential and regional expression of mRNA for basic fibroblast

growth factor in the developing and

adult rat brain, 71

Printz, D., see Angulo, J.A., 301

Rius, R.A., see Wong, M., 197 Roberson, M.D., see Toews, A.D., 321 Rodríguez, E.M., see Nualart, F., 227

Sabban, E.L., see Badoyannis, H.C., 79 Saika, T., Senba, E., Noguchi, K., Sato, M., Kubo, T., Matsunaga, T. and Tohyama, M., Changes in expression of peptides in rat facial motoneurons after facial nerve crushing and resection, 187

Sato, M., see Saika, T., 187 Sato, M., see Zhang, J.-H., 239 Scarth, B.A., see Shaw, C., 273 Scheller, R.H., see Maroteaux, L., 335 Scheller, R.H., see Volknandt, W., 283 Schilling, J., see Volknandt, W., 283 Schlichter, D., see Huang, Z., 309

Schreiber, S.S., Maren, S., Tocco, G., Shors, T.J. and Thompson, R.F., A negative correlation between the induction of long-term potentiation and activation of immediate early genes, 89 Schumacher, M., see Angulo, J.A., 1

Schwartz, M., see Cohen, I., 181 Schwartz, J.P., see Shinoda, H., 221 Seitanidou, T., see Becker, C.-M., 327 Senba, E., see Saika, T., 187 Shani, Y., see Cohen, I., 181 Shanker, G., see Claro, E., 265

Sharma, S.C., see Badoyannis, H.C., 79 Shaw, C. and Scarth, B.A., Characterization and differential regulation of GABA_A and benzodiazepine receptors in rat neocortex, 273

Shin, C., see Simonato, M.,115

Shinoda, H., Nadi, N.S. and Schwartz, J.P., Alterations in somatostatin and proenkephalin mRNA in response to a single amygdaloid stimulation versus kindling, 221

Shors, T.J., see Schreiber, S.S., 89 Simonato, M., Hosford, D.A., Labiner, D.M., Shin, C., Mansbach, H.M. and McNamara, J.O., Differential expression of immediate early genes in the hippocampus in the kindling model of epilepsy,115

Sirinathsinghji, D.J.S., see Matthews, S.G., 171

Snutch, T.P., see Yu, L., 143 Stone, E.A., see Bing, G., 43 Streamson C. Chua Jr., Andrew W. Brown, Jeehee Kim, Karen L. Hennessey, Rudolph L. Leibel and Hirsch, J., Food deprivation and hypothalamic neuropeptide gene expression: effects of strain background diabetes and the mutation, 291

Sudol, M., see Lustig, R.H., 125 Sugar, J., see Perez, N., 249 Sumikawa, K., see Gehle, V.M., 17

Teskey, G.C., Atkinson, B.G. and Cain, D.P., Expression of the proto-oncogene c-fos following electrical kindling in the rat, 1

Thewke, D., see Huang, Z., 309 Thompson, R.F., see Schreiber, S.S., 89 Tocco, G., see Schreiber, S.S., 89

Toews, A.D., Eckermann, C.E., Roberson, M.D., Lee, S.Y. and Morell, P., Primary demyelination induced by exposure to tellurium alters mRNA levels for nerve growth factor receptor, SCIP, 2',3'-cyclic nucleotide 3'-phosphodiesterase, and myelin proteolipid protein in rat sciatic nerve, 321

Tohyama, M., see Saika, T., 187 Tohyama, M., see Zhang, J.-H., 239 Triller, A., see Becker, C.-M., 327 Turk, C.W., see Mitsuhashi, M., 177

Uhl, G.R., Walther, D., Nishimori, T., Buzzi, M.G. and Moskowitz, M.A., Jun B, c-jun, jun D and c-fos mRNAs in nucleus caudalis neurons: rapid selective enhancement by afferent stimulation, 133

Van Heerikhuizen, H., see Bogerd, J., 47 Volknandt, W., Pevsner, J., Elferink, L.A., Schilling, J. and Scheller, R.H., A synaptic vesicle specific GTP-binding protein from ray electric organ, 283

Wallace, W., see Perez, N., 249 Wallace, M.A., see Claro, E., 265 Walther, D., see Uhl, G.R., 133 Weiss, S.R.B., see Clark, M., 55

Whiting, P., Nava, S., Mozley, L., Eastham, H. and Poat, J., Expression of angiotensin converting enzyme mRNA in rat brain, 1

Wicks, W.D., see Huang, Z., 309

Wong, M., Rius, R.A. and Loh, Y.P., Characterization of Xenopus laevis proenkephalin gene, 197

Yu, L., Nguyen, H., Le, H., Bloem, L.J., Kozak, C.A., Hoffman, B.J., Snutch, T.P., Lester, H.A., Davidson, N., and Lübbert, H., The mouse 5-HT_{1C} receptor contains eight hydrophobic domains and is X-linked, 143

Zhang, J.-H., Araki, T., Sato, M. and Tohyama, M., Distribution of GABAAreceptor a_1 subunit gene expression in the rat forebrain, 239

